

BRAIDING KNOTS IN CONTACT 3-MANIFOLDS

ELENA PAVELESCU

ABSTRACT. We show that a transverse link in a contact structure supported by an open book decomposition can be transversely braided. We also generalize Markov's theorem on when the closures of two braids represent (transversely) isotopic links.

1. INTRODUCTION

In [1], Alexander proved that any knot in $\mathbb{R}^3$ can be braided about the z -axis. In [3], a paper which marked the start of modern contact topology, leading to the Bennequin inequality and eventually to Eliashberg's definition of tightness, Bennequin proved the transverse case for $(\mathbb{R}^3, \xi_{std})$. In this paper, we generalize Bennequin's result to any closed oriented 3-dimensional manifold M , by looking at an open book decomposition for M together with its supported contact structure. After completing the proof of this theorem, it was brought to our attention that the same result was independently obtained by Mitsumatsu and Mori in [8].

Bennequin used braid theory in $\mathbb{R}^3$ to show the standard contact structure was tight. Since then, the main way to show a contact structure is tight is to use powerful analytic machinery like holomorphic curves or Seiberg-Witten theory. By studying braids in general open books it is hoped that tightness can be better understood from a purely topological/combinatorial perspective. The results presented in this paper constitute a first step in this program. While these results are purely geometric, it would be nice to have an algebraic description of a generalized "braid group". In particular, one needs to define the stabilization moves with respect to different binding components and account for the fundamental group of the page. We will explore this algebraic perspective in a subsequent paper.

In [7], Markov gave an equivalent condition for the closures of two braids in $\mathbb{R}^3$ to be isotopic as knots. This is the case if and only if the two braids differ by conjugations in the braid group and positive and negative Markov moves. In [9], Orevkov and Shevchishin proved the transversal case for $(\mathbb{R}^3, \xi_{std})$. A different proof was independently obtained by Wrinkle in [13]. We generalize Markov's theorem to any closed oriented 3-dimensional manifold. We prove the transverse case and recover the topological case previously proved in [10] and [11].

ACKNOWLEDGEMENTS

The author would like to thank John Etnyre for all his support and guidance during the past years, and for all the valuable comments and suggestions about this paper. She would also like to thank Keiko Kawamuro for all the helpful conversations.

2. CONTACT STRUCTURES AND OPEN BOOK DECOMPOSITIONS

Let M be a compact, oriented 3-manifold, and ξ an oriented 2-plane field on M . We say ξ is a *contact structure* on M if $\xi = \ker \alpha$ for some global 1-form α satisfying $\alpha \wedge d\alpha \neq 0$. In this paper we assume that the manifold M is oriented and that the contact structure ξ is oriented and positive (i.e. $\alpha \wedge d\alpha > 0$).

A *contactomorphism* between two contact manifolds (M_1, ξ_1) and (M_2, ξ_2) is a diffeomorphism $\phi : M_1 \rightarrow M_2$ such that $\phi_*\xi_1 = \xi_2$.

On $\mathbb{R}^3$, consider the two contact structures ξ_1 and ξ_2 given by the 1-forms $\alpha_1 = dz - ydx$ and $\alpha_2 = dz + r^2 d\theta$ (given in cylindrical coordinates). Then $(\mathbb{R}^3, \xi_1)$ and $(\mathbb{R}^3, \xi_2)$ are contactomorphic and we are going to refer to both ξ_1 and ξ_2 as the standard contact structure on $\mathbb{R}^3$, ξ_{std} .

Theorem 2.1. (Gray) *Let $\{\xi_t\}_{t \in [0,1]}$ be a family of contact structures on a manifold M that differ on a compact set $C \subset \text{int}(M)$. Then there exists an isotopy $\psi_t : M \rightarrow M$ such that*

- i) $(\psi_t)_*\xi_1 = \xi_t$
- ii) ψ_t is the identity outside of an open neighborhood of C .

While the proof of this result is well known, we sketch it here as we will need elements of it in later arguments. For more details see [5] (p. 61).

Proof. We are going to look for ψ_t as the flow of a vector field X_t . If $\xi_t = \ker \alpha_t$, then ψ_t has to satisfy

$$(2.1) \quad \psi_t^* \alpha_t = \lambda_t \alpha_0,$$

for some non-vanishing function $\lambda_t : M \rightarrow \mathbb{R}$. By taking the derivative with respect to t on both sides and rearranging the terms we get

$$(2.2) \quad \psi_t^* \left(\frac{d\alpha_t}{dt} + \mathcal{L}_{X_t} \alpha_t \right) = \frac{d\lambda_t}{dt} \alpha_0 = \frac{d\lambda_t}{dt} \frac{1}{\lambda_t} \psi_t^* \alpha_t.$$

This is equivalent to

$$(2.3) \quad \psi_t^* \left(\frac{d\alpha_t}{dt} + d(\iota_{X_t} \alpha_t) + \iota_{X_t} d\alpha_t \right) = \psi_t^* (h_t \alpha_t)$$

for

$$h_t = \frac{d}{dt} (\log \lambda_t) \circ \psi_t^{-1}.$$

If X_t is chosen in ξ_t then $\iota_{X_t} \alpha_t = 0$ and (2.3) becomes

$$(2.4) \quad \frac{d\alpha_t}{dt} + \iota_{X_t} d\alpha_t = h_t \alpha_t$$

Applying (2.4) to the Reeb vector field of α_t , v_{α_t} (that is, the unique vector field such that $\alpha_t(v_t) = 1$ and $d\alpha_t(v_t, \cdot) = 0$), we find $h_t = \frac{d\alpha_t}{dt}(v_{\alpha_t})$ and X_t given by

$$(2.5) \quad \iota_{X_t} d\alpha_t = h_t \alpha_t - \frac{d\alpha_t}{dt}$$

The form $d\alpha_t$ gives an isomorphism

$$\begin{aligned}\Gamma(\xi_t) &\rightarrow \Omega_{\alpha_t}^1 \\ v &\mapsto \iota_v d\alpha_t\end{aligned}$$

where $\Gamma(\xi_t) = \{v | v \in \xi_t\}$ and $\Omega_{\alpha_t}^1 = \{1\text{-forms } \beta | \beta(v_t) = 0\}$, and thus X_t is uniquely determined by (2.4). By construction, the flow of X_t works. For the subset of M where the ξ_t 's agree we just choose the α_t 's to agree. This implies $\frac{d\alpha_t}{dt} = 0$, $h_t = 0$ and $X_t = 0$ and all equalities hold. $\square$

In a contact manifold (M, ξ) , an oriented arc $\gamma \subset M$ is called *transverse* if for all $p \in \gamma$ and ξ_p the contact plane at p , $T_p \gamma \pitchfork \xi_p$ and $T_p \gamma$ intersects ξ_p positively. If γ is a closed curve then it is called a *transverse knot*.

An *open book decomposition* of M is a pair (L, π) where

- (1) L is an oriented link in M called *the binding* of the open book
- (2) $\pi : M \setminus L \rightarrow S^1$ is a fibration whose fiber, $\pi^{-1}(\theta)$, is the interior of a compact surface $\Sigma \subset M$ such that $\partial \Sigma = L$, for all $\theta \in S^1$. The surface Σ is called *the page* of the open book.

Alternatively, an open book decomposition of a 3-manifold M consists of a surface Σ , with boundary, together with a diffeomorphism $\phi : \Sigma \rightarrow \Sigma$, with ϕ the identity near $\partial \Sigma$, such that

$$M = (\Sigma \times [0, 1] / \sim) \cup_f \coprod_i S^1 \times D^2$$

where

$$(x, 1) \sim (\phi(x), 0).$$

Note that

$$\partial(\Sigma \times [0, 1] / \sim) = \coprod_i T_i^2,$$

with each torus T_i^2 having a product structure $S^1 \times [0, 1] / \sim$. Let $\lambda_i = \{pt\} \times [0, 1] / \sim$, $\lambda_i \in T_i^2$. The gluing diffeomorphism used to construct M is defined by

$$\begin{aligned}f : \partial(\coprod_i S^1 \times D^2) &\rightarrow \partial(\coprod_i T_i^2) \\ \{pt\} \times \partial D^2 &\rightarrow \lambda_i.\end{aligned}$$

The map ϕ is called *the monodromy* of the open book. See [4] for more details.

Theorem 2.2. (Alexander, [2]) *Every closed oriented 3-manifold has an open book decomposition.*

A contact structure ξ on M is said to be *supported* by an open book decomposition (Σ, ϕ) of M if ξ can be isotoped through contact structures so that there exists a 1-form α for ξ such that

- (1) $d\alpha$ is a positive area form on each page
- (2) $\alpha(v) > 0$ for all $v \in TL$ that induce the orientation on L .

Theorem 2.3. (Thurston, Winkelnkemper, [12]) *Every open book decomposition (Σ, ϕ) supports a contact structure ξ_ϕ .*

We sketch this well know proof as we need the details in later arguments.

Proof. Let

$$M = (\Sigma \times [0, 1] / \sim) \cup_f \coprod_i S^1 \times D^2$$

given as before. We first construct a contact structure on $\Sigma \times [0, 1] / \sim$ and then we extended it in a neighborhood of the binding.

In the neighborhood $N = S^1 \times D^2$ of each component of the binding consider coordinates (ψ, x, θ) such that (ψ, x) are coordinates on the page with ψ the coordinate along the binding and $d\theta$ and $\pi^*d\theta$ agree, where $\pi^*d\theta$ is the pullback through $\pi : M \setminus L \rightarrow S^1$ of the coordinate on S^1 . Let λ be a 1-form on the page which is an element of the set (it is easy to check that this set is not empty)

$$S = \{1\text{-forms } \lambda \text{ such that } : d\lambda \text{ is a volume form on } \Sigma \\ \lambda = (1 + x)d\psi \text{ near } \partial\Sigma = L\}$$

On $\Sigma \times [0, 1]$ take $\tilde{\lambda} = (1 - \theta)\lambda + \theta(\phi^*\lambda)$ and consider the 1-form $\alpha = \tilde{\lambda} + Kd\theta$. For sufficiently large K , α_K is a contact form and it descends to a contact form on $\Sigma \times [0, 1] / \sim$. To extend this form on the solid tori neighborhood of the binding we pull back α through the gluing map f and get

$$\alpha_f = Kd\theta - (r + \epsilon)d\psi.$$




FIGURE 1. h and g functions

We are looking to extend this form on the entire $S^1 \times D^2$ to a contact form of the form $h(r)d\psi + g(r)d\theta$. This is possible if there exist functions $h, g : [0, 1] \rightarrow \mathbb{R}^3$ such that:

- (1) $h(r)g'(r) - h'(r)g(r) > 0$ (given by the contact condition)
- (2) $h(r) = 1$ near $r = 0$, $h(r) = -(r + \epsilon)$ near $r = 1$
- (3) $g(r) = r^2$ near $r = 0$, $g(r) = K$ near $r = 1$.

The two functions h and g described in Figure 1 work for our purpose. The conditions (1) and (2) are obviously satisfied, and if δ_h and δ_g are such that $h(r) < 0$ on $[\delta_h, 1]$ and $g(r) = 1$ on $[\delta_g, 1]$, then (3) is satisfied as long as $\delta_h < \delta_g$.

□

The following correspondence, which is one of the central results in contact geometry, relates open book decompositions and contact structures.

Theorem 2.4. (*Giroux, [6]*) *Let M be a closed, oriented 3-manifold. Then there is a one to one correspondence*

$$\left\{ \begin{array}{l} \text{Contact structures } \xi \text{ on } M^3 \\ \text{up to contact isotopy} \end{array} \right\} \xleftrightarrow{1-1} \left\{ \begin{array}{l} \text{Open book decompositions } (\Sigma, \phi) \\ \text{of } M^3 \text{ up to positive stabilization} \end{array} \right\}.$$

See [4] for details.

3. BRAIDING KNOTS IN 3-MANIFOLDS

In this section we generalize the following theorem:

Theorem 3.1. (*Bennequin, [3]*) *Any transverse link K in $(\mathbb{R}^3, \xi_{std})$ is transversely isotopic to a link braided about the z -axis.*

Let (L, π) be an open book decomposition for M . A link $K \subset M$ is said to be *braided about L* if K is disjoint from L and there exists a parametrization of K , $f : \coprod S^1 \rightarrow M$, such that if θ is the coordinate on each S^1 then $\frac{d}{d\theta}(\pi \circ f) > 0, \forall \theta$. We call *bad arcs* of K those arcs where this condition is not satisfied.

Theorem 3.2. *Suppose (L, π) is an open book decomposition for the 3-manifold M and ξ is supported by (L, π) . Let K be a transverse link in M . Then K can be transversely isotoped to a braid.*

Proof. The idea of the proof is to find a family of diffeomorphisms of M keeping each page of the open book setwise fixed and taking the parts of the link where the link is not braided in a neighborhood of the binding. A neighborhood of the binding is contactomorphic to a neighborhood of the z -axis in $(\mathbb{R}^3, \xi_{std})/_{z \sim z+1}$ and there the link can be braided, according to Theorem 3.1. What we really use is that Theorem 3.1 works not only for links, but also for arcs with good ends.

In the neighborhood $N = S^1 \times D^2$ of each component of the binding consider coordinates (ψ, x, θ) and let $\lambda \in S$ as described in Theorem 2.3. On $\Sigma \times [0, 1]$ take $\tilde{\lambda} = (1 - \theta)\lambda + \theta(\phi^*\lambda)$ and consider the family of 1-forms given by $\alpha_t = \tilde{\lambda} + K \frac{1}{t} d\theta$, where $t \in (0, 1]$ and K is any large constant.

This family of 1-forms descends to a family of 1-forms on $\Sigma \times [0, 1]/\sim$ (because $\phi^*(\alpha_t|_{\Sigma \times \{0\}}) = \alpha_t|_{\Sigma \times \{1\}}$). Both $\xi_1 = \ker(\alpha_1)$ and ξ are contact structures supported by (L, π) and by Giroux's correspondence they are isotopic. Therefore, without loss of generality, we may assume $\xi = \ker(\alpha_1)$.

For large enough K , the family of 1-forms $\{\alpha_t\}_t$ is a family of contact 1-forms as:

$$\alpha_t \wedge d\alpha_t = (\tilde{\lambda} + K \frac{1}{t} d\theta) \wedge (d\tilde{\lambda}) = \tilde{\lambda} \wedge d\tilde{\lambda} + K \frac{1}{t} d\theta \wedge d\tilde{\lambda} > 0$$

Note that $d\tilde{\lambda}$ is an area form on the page while $d\theta$ vanishes on the page and is positive on the positive normal to the page. This implies that the second term of the sum is always positive and therefore α_t is a contact form for sufficiently large K . We want to extend this family to the whole M , so we need to patch in the solid tori neighborhood of the binding. In the neighborhood of the binding we have $\alpha_t = (1 + x)d\psi + K \frac{1}{t} d\theta$. We

pull back this family through the map f used to glue the solid tori in the definition of the open book. With our chosen coordinates we have

$$f(\psi, x, \theta) = (x - 1 + \epsilon, -\psi, \theta)$$

and the pullback family

$$\alpha_{f,t} = K \frac{1}{t} d\theta - (x + \epsilon) d\psi.$$

We are looking to extend this forms on the entire $S^1 \times D^2$ to a family of the form $h_t(x)d\psi + g_t(x)d\theta$. This is possible if there exist functions $h_t, g_t : [0, 1] \rightarrow \mathbb{R}$, for $t \in (0, 1]$ such that:

- i) $h_t(x)g'_t(x) - h'_t(x)g_t(x) > 0, t \in (0, 1]$ (given by the contact condition)
- ii) $h_t(x) = 1$ near $x = 0$, $h_t(x) = -(x + \epsilon)$ near $x = 1$
- iii) $g_t(x) = x^2$ near $x = 0$, $g_t(x) = \frac{K}{t}$ near $x = 1$.




FIGURE 2. h_t and g_t functions

The two families $\{h_t\}_{t \in (0,1]}$ and $\{g_t\}_{t \in (0,1]}$ described in Figure 2 work for our purpose. The conditions i) and ii) are obviously satisfied and if δ_{h_t} and δ_{g_t} are such that $h_t(x) < 0$ for $x \in [\delta_{h_t}, 1]$ and $g_t(x) = 1$ for $x \in [\delta_{g_t}, 1]$, then iii) is satisfied as long as $\delta_{h_t} < \delta_{g_t}$.

Denote the extended family of forms also by α_t and by ξ_t the family of contact structures given by $\xi_t = \ker(\alpha_t)$, $t \in (0, 1]$. By Gray's theorem there exists a family of diffeomorphisms $f_t : M \rightarrow M$ such that $(f_t)_*\xi_1 = \xi_t$.

As announced, we would like to construct a family of diffeomorphisms $\{f_t\}_t$ that fix the pages setwise. Following the proof of Theorem 2.1, we construct $\{f_t\}_t$ as the flow of a vector field $X_t \in \xi_t$, for which we have the equality (2.5).

We already know that such a X_t exists but would need it to be tangent to the page. First notice that $\frac{d\alpha_t}{dt} = -\frac{1}{t^2}Kd\theta$ and choose some vector $v \in T\Sigma \cap \xi_t$. Applying both sides of (2.5) to v we get

$$(3.1) \quad d\alpha_t(X_t, v) = \frac{d\alpha_t}{dt}(v)\alpha_t(v) - \frac{d\alpha_t}{dt}(v)$$

As $v \in \xi_t = \ker(\alpha_t)$ and v has no θ -component, the last equality is equivalent to

$$(3.2) \quad d\alpha_t(X_t, v) = 0.$$

As $d\alpha_t$ is an area form on ξ_t , this implies that X_t and v are linearly dependent and therefore $X_t \in T\Sigma \cap \xi_t$ ($X_t = 0$ at singular points, where $T\Sigma$ and ξ_t coincide).

We are now looking at the singularities of $X_t, t \in (0, 1]$. Because for $t_1 \neq t_2$ α_{t_1} and α_{t_2} differ by a multiple of $d\theta$, the flowlines given by X_{t_1} and X_{t_2} on a page Σ_θ coincide. Therefore, it suffices to look at the singularities of X_{t_0} for some $t_0 \in (0, 1]$. On Σ_θ there are no negative elliptic singularities away from the binding as the contact planes and the planes tangent to the page almost coincide, as oriented plane fields (a negative elliptic singularity e would require ξ_e and $T_e\Sigma$ to coincide but have different orientations). Thus, for each θ , all points on Σ_θ , except for singularities of X_t and stable submanifolds of hyperbolic points, flow in finite time into an arbitrarily small neighborhood of the binding. Define S_θ as the set of points on Σ_θ that are either singularities of X_t or on stable submanifolds of hyperbolic points. Let $\mathcal{S} = \cup S_\theta$ as θ varies from 0 to 2π .

First, note that we can arrange the monodromy map ϕ to fix the singularities on the cutting page, by thinking of ϕ as of a composition of Dehn twists away from these points. For isolated values of θ , X_t might exhibit connections between hyperbolic singularities. With these said, $\mathcal{S}$ has a *CW* structure with

1-skeleton: union of $\{x\} \times [0, 1]$ for singular points x , and connections between hyperbolic singularities

2-skeleton: union of stable submanifolds of hyperbolic singularities.

We want to arrange K in such a way that the bad arcs of K (where the knot is not braided) are disjoint from $\mathcal{S}$. We introduce wrinkles along K as in Figure 2. See below for the explicit definition.




FIGURE 3. Wrinkling K in order to avoid intersections with $\mathcal{S}$

The wrinkles may increase the number of arcs where the link is not braided but this is fine, as these new arcs avoid $\mathcal{S}$.

By general position, we may assume $K \cap (1 - \text{skeleton of } S) = \emptyset$ and $K \cap (2 - \text{skeleton of } S)$ is a finite number of points. A small neighborhood of $p \in K \cap (2 - \text{skeleton of } S)$ in Σ_θ is foliated by intervals $(-\epsilon, \epsilon)$, in the same way as a small disk in the xy -plane centered at $(0, 1, 0)$ in $(\mathbb{R}^3, \xi_{std})$. It follows that p has a neighborhood in M which is contactomorphic to a neighborhood of $q = (0, 1, 0)$ in $(\mathbb{R}^3, \xi_{std})$. Consider the standard (x, y, z) coordinate system in such a neighborhood of q . The contact plane at q is given by the equation $z = x$. As at p the contact plane and the plane tangent to the page almost coincide, we consider the plane $z = (1 + \epsilon)x$ at q to correspond to the plane tangent to the page at p . In a δ -neighborhood of q in $\mathbb{R}^3$, the line given by $y = 1, z = (1 + \frac{\epsilon}{2})x$ will represent the corresponding arc of K . With this setting, the wrinkle takes K to K' with the following properties:

i) K' is given by $z = 1 + \frac{3\epsilon}{2}x, y = 1$ in a $\frac{\delta}{3}$ -neighborhood of q

- ii) K' is given by $z = 1 + \frac{\epsilon}{2}x$, $y = 1$ outside of a $\frac{2\delta}{3}$ -neighborhood of q
- iii) $\frac{dz}{dx} > 0$ along K' in a δ -neighborhood of q .

While condition i) takes care of the K avoiding S along its bad arcs, condition iii) takes care of the link remaining transverse throughout the wrinkling.

After making the necessary wrinkles and applying the family of diffeomorphisms $\{f_t\}_{t \in (0,1]}$, $K_0 := f_\epsilon(K)$ has all bad regions in a neighborhood of the binding and by Theorem 3.1 there is a transverse isotopy K_s , $0 \leq s \leq 1$ taking K_0 to a braid K_1 . The wrinkling isotopy followed by K_s is the transverse isotopy we are looking for. $\square$

4. MARKOV'S THEOREM IN AN OPEN BOOK DECOMPOSITION

The goal of this section is to generalize the following theorem

Theorem 4.1. (*Orevkov, Shevchishin [9]*) *In $(\mathbb{R}^3, \xi_{std})$ two braids represent transversely isotopic links if and only if one can pass from one braid to the other by braid isotopies, positive Markov moves and their inverses.*

A *Markov move* near a component of the binding L is given by the introduction of a new loop around L , as in Figure 3. These moves are done near the binding, where the standard coordinates apply. While a positive Markov move (and its inverse) keeps the link transverse, a negative Markov move (and its inverse) fail to do so (see [3] for details).



FIGURE 4. Positive (left) and negative (right) Markov moves.

Theorem 4.2. *Let M be a 3-dimensional, closed, oriented manifold and (L, π) an open book decomposition for M together with its supported contact structure ξ . Let K_0 and K_1 be transverse closed braid representatives of the same topological link. Then K_0 and K_1 are transversely isotopic if and only if they differ by braid isotopies and positive Markov moves and their inverses.*

The topological version has been previously proven by Skora [10] and Sundheim [11]. This case immediately follows from the proof of the transverse case, as one does not need to worry about transversality throughout the isotopy and if transversality is not required both positive and negative Markov moves are allowed.

Proof. The reverse implication is straightforward. We only need to note that an isotopy through braids is done away from the binding. As the contact planes almost coincide to the planes tangent to the pages this isotopy stays transverse with respect to the contact structure.

The direct implication takes more work. Let K_0 and K_1 be transverse braid representatives of the same topological link K and $\{K_t\}_{t \in [0,1]}$ a transverse isotopy from K_0 to K_1 . We parametrize the isotopy by $\mathcal{K} : \sqcup S^1 \times [0, 1] \rightarrow M$, such that $\mathcal{K}_t$ defined by $s \rightarrow \mathcal{K}(s, t)$ is a parametrization of K_t , where s is the positively oriented coordinate on each S^1 .




FIGURE 5. Wrinkling in order to avoid intersections between $\mathcal{B}$ and $\mathcal{S}$

Let θ be the positive coordinate normal to the page. A *bad zone* of $\mathcal{K}$ is a connected component of the set of points in $\sqcup S^1 \times [0, 1]$ for which $\frac{\partial(\pi \circ \mathcal{K})}{\partial s} \leq 0$. Denote by $\mathcal{B}$ the union of all bad zones of $\mathcal{K}$.

We would like to take all the bad zones of $\mathcal{K}$ in a neighborhood of the binding. This way the proof is reduced to the standard case proved by Orevkov and Shevchishin in [9]. For this, we use the family of diffeomorphisms $\{f_t\}_{t \in (0,1]}$ constructed in the proof of Theorem 3.2. Keeping the same notations, we want to arrange the isotopy $\mathcal{K}$ in such a way that $\mathcal{B} \cap \mathcal{S} = \emptyset$. By general position, $\mathcal{B} \cap \mathcal{S}$ consists of arcs l , along which $\frac{\partial(\pi \circ \mathcal{K})}{\partial s} \leq 0$ and points at which $\frac{\partial(\pi \circ \mathcal{K})}{\partial s} \leq 0$. The arcs l are intersections of the 2-cells of $\mathcal{S}$ with $\mathcal{B}$ while the points are intersections of the 1-cells of $\mathcal{S}$ with $\mathcal{B}$. We modify the isotopy $\mathcal{K}$ in such a way so as to replace the arc $l \subset \mathcal{B} \cap \mathcal{S}$ by another arc $l' \subset \mathcal{S}$, along which $\frac{\partial(\pi \circ \mathcal{K})}{\partial s} > 0$ and to eliminate the isolated points of $\mathcal{B} \cap \mathcal{S}$. We describe the process below.

By general position, we may assume that the isotopy has at most triple points and that $\mathcal{S}$ does not intersect $\mathcal{B}$ in one of these triple points. Let $x \in l \subset \mathcal{B} \cap \mathcal{S}$. The point x can be at most a double point of the immersed annulus.

Case 1. If x is not a double point, then pick coordinates (s, t) in a small neighborhood of x on the isotopy annulus (where s is the coordinate along S^1 and t is the coordinate along the interval factor, $[0, 1]$). Assume that $x = (s_0, t_0)$ in these coordinates. In this neighborhood, we modify $\mathcal{K}((s_0 - \epsilon, s_0 + \epsilon) \times \{t_0\})$ for a small $\epsilon > 0$ as described in Theorem 3.2, and we do this continuously with respect to the t -coordinate. For all points $x \in l$, we have $x = (p_x, \theta_x) \in \Sigma_{\theta_x} \times \{\theta_x\}$. We keep the page coordinate p_x fixed and we modify the θ -coordinate as in Figure 5. As the manifold is compact, l may be covered with finitely many such neighborhoods and after finitely many such modifications $\mathcal{B} \cap \mathcal{S}$ has one less arc. See Figure 6.

Case 2. If x is a double point on the immersed annulus, then we can find two different neighborhoods of x on the annulus, $U_1 \subset \mathcal{K}((s_0, s_1) \times (t_0, t_1))$ and $U_2 \subset \mathcal{K}((s_2, s_3) \times (t_2, t_3))$, with $(s_0, s_1) \cap (s_2, s_3) = \emptyset$ and $(t_0, t_1) \cap (t_2, t_3) = \emptyset$. We look at one of these




FIGURE 6. Isotopy modification along an l arc. The new intersection arc sits outside of $\mathcal{B}$

neighborhoods, say U_1 , and modify the isotopy, as described above, only for U_1 and then continuously with respect to the t -coordinate along neighborhoods of points on l included in the same branch as U_1 . In this way we reduce the proof to the previous case.

If x is an isolated point of $\mathcal{S} \cap \mathcal{B}$ we need only modify the isotopy in a small neighborhood of x . After finitely many such modifications, $\mathcal{B} \cap \mathcal{S} = \emptyset$. $\square$

REFERENCES

- [1] J.W.Alexander, *A lemma on systems of knotted curves*, Proc. Nat. Acad. Sci. USA, **9** (1923), 93-95
- [2] J.W. Alexander, *Note on Riemann spaces*, Bull. Amer. Math. Soc., **26** (1920), 370-372
- [3] D.Bennequin, *Entrelacements et équations de Pfaff*, Astérisque **107-108** (1983), 87-161
- [4] J.Etnyre, *Lectures on open book decompositions and contact structures*, Clay Mathematics Proceedings, Volume **5** (2006), 103-142
- [5] H. Geiges, *An Introduction to Contact Topology*, Cambridge Studies in Advanced Mathematics, No. **109**, (2008)
- [6] E.Giroux, *Géométrie de contact: de la dimension trois vers les dimensions supérieures*, Proceedings of the International Congress of Mathematicians, Vol. II, 405-414, Higher Ed. Press, Beijing, (2002)
- [7] A.A. Markov, *Über die freie Äquivalenz der geschlossenen Zöpfe*, Mat. Sbornik **43** (1936), 73-78
- [8] Y. Mitsumatsu, *Convergence of contact structures to foliations. With an appendix by Mitsumatsu and Atsuhide Mori*. Foliations 2005, 353-371, World Sci. Publ., Hackensack, NJ, 2006.
- [9] S.Yu.Orevkov, V.V.Shevchishin, *Markov theorem for transversal links*, J.of Knot Theory and its Ramifications, **12** (2003), 905-913
- [10] R.Skora, *Closed braids in 3-manifolds*, Mathematische Zeitschrift, Volume **211**, No.1 (1992), 173-187
- [11] P.Sundheim, *The Alexander and Markov Theorems via Diagrams for Links in 3-Manifolds*, Trans. Amer. Math. Soc., **2** (1993), 591-607
- [12] W.P.Thurston, H.Winkelkemper, *On the Existence of Contact Forms*, Proc. Amer. Math. Soc. **52** (1975), 345-347
- [13] N. Wrinkle, *The Markov theorem for transverse knots*, arXiv:math/0202055v1

RICE UNIVERSITY, DEPARTMENT OF MATHEMATICS - MS 136, HOUSTON, TX 77005
E-mail address: Elena.Pavelescu@rice.edu